

Econometrics

Chapter 18: Cointegration

In Choi

Sogang University

- A series with no deterministic component which has a stationary, invertible, ARMA representation after differencing d times, is said to be integrated of order d , denoted $x_t \sim I(d)$.

Example

$$y_t = y_{t-1} + u_t, u_t \sim WN(0, \sigma^2).$$

Then $y_t \sim I(1)$.

- Properties of $I(1)$ series

(i) Growing variance. ($y_t \sim I(1)$. Then $\text{var}(y_t) \rightarrow \infty$ as $t \rightarrow \infty$.)

(ii) An innovation has a permanent effect on the value of y_t .

$$(y_t = \sum_{-\infty}^t u_i.)$$

(iii) The expected time between crossings of $x = 0$ is infinite.

(iv) The theoretical autocorrelation, $\rho_k \rightarrow 1$ for all k as $t \rightarrow \infty$.

- If $x_t \sim I(1)$ and $y_t \sim I(1)$, it is generally true that $z_t = y_t - ax_t \sim I(1)$. When $z_t \sim I(0)$, we say that x_t and y_t are cointegrated.

Example

- (i) x_t : dividend, y_t : stock price
- (ii) x_t : income, y_t : consumption
- (iii) x_t : interest rate, y_t : money

- The coefficient a represents the long-run equilibrium relationship among variables.
- Consider the regression model

$$y_t = \mu + \alpha x_t + u_t, \quad (t = 1, \dots, T)$$

where $x_t \sim I(1)$ and $u_t \sim I(0)$.

- Even if Δx_t and u_t are correlated,

$$\hat{\alpha} \xrightarrow{P} \alpha.$$

- $T(\hat{\alpha} - \alpha)$ has a nonnormal limiting distribution. This is different from the usual \sqrt{T} -asymptotics.

- There are various methods for estimating α efficiently.
- How can we know x and y are cointegrated?
Step 1: Run OLS and get residuals $\{\hat{u}_t\}$.
Step 2: Test for a unit root using $\{\hat{u}_t\}$. (If unit root, non-cointegration. If no unit root, cointegration.)
- More sophisticated methods use the vector autoregression in order to test for the presence of cointegration.